



**Biomass, Size, and Trophic Status of Top Predators
in the Pacific Ocean**

John Sibert, *et al.*
Science **314**, 1773 (2006);
DOI: 10.1126/science.1135347

***The following resources related to this article are available online at
www.sciencemag.org (this information is current as of December 14, 2006):***

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/314/5806/1773>

Supporting Online Material can be found at:

<http://www.sciencemag.org/cgi/content/full/314/5806/1773/DC1>

This article **cites 5 articles**, 3 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/314/5806/1773#otherarticles>

This article appears in the following **subject collections**:

Ecology

<http://www.sciencemag.org/cgi/collection/ecology>

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/help/about/permissions.dtl>

Biomass, Size, and Trophic Status of Top Predators in the Pacific Ocean

John Sibert,^{1*} John Hampton,² Pierre Kleiber,³ Mark Maunder⁴

Fisheries have removed at least 50 million tons of tuna and other top-level predators from the Pacific Ocean pelagic ecosystem since 1950, leading to concerns about a catastrophic reduction in population biomass and the collapse of oceanic food chains. We analyzed all available data from Pacific tuna fisheries for 1950–2004 to provide comprehensive estimates of fishery impacts on population biomass and size structure. Current biomass ranges among species from 36 to 91% of the biomass predicted in the absence of fishing, a level consistent with or higher than standard fisheries management targets. Fish larger than 175 centimeters fork length have decreased from 5% to approximately 1% of the total population. The trophic level of the catch has decreased slightly, but there is no detectable decrease in the trophic level of the population. These results indicate substantial, though not catastrophic, impacts of fisheries on these top-level predators and minor impacts on the ecosystem in the Pacific Ocean.

Industrial fisheries for tunas and associated species extend over most of the tropical and temperate Pacific Ocean and currently produce over 2.5 million tons (*I*) or about 64% of the 2004 global tuna catch (Fig. 1). These fisheries have grown continuously since the 1950s and have removed more than 50 million tons of large pelagic fish. The fishery targets four primary temperate and tropical tuna species: skipjack (*Katsuwonus pelamis*), yellowfin (*Thunnus albacares*), bigeye (*T. obesus*), and albacore (*T. alalunga*) tuna. Other species, blue-

fin tuna (*T. orientalis*), billfishes, and oceanic sharks are also taken, but the primary tuna species make up >90% of the total catch by weight. Data derived from these fisheries (including catch, fishing effort, size composition, and tagging data) show a 50-year record of natural variability in and human impacts on open-ocean ecosystems. We analyzed all available data with state-of-the-art stock assessment methods to provide estimates of fishery impacts on population biomass, size structure, and trophic status of major top-level predator stocks in the Pacific

Ocean: bigeye tuna, yellowfin tuna, skipjack tuna, albacore tuna, and blue shark (*Prionace glauca*) (2).

The trajectories of exploited and unexploited (3) biomass vary substantially among stocks (Fig. 2). Exploited western Pacific yellowfin and bigeye have declined steadily to levels near the equilibrium biomass that would produce the maximum sustainable yield (MSY) in the fishery. Skipjack tuna and blue shark appear to have increased slightly, whereas albacore have fluctuated in both directions. Current total and adult biomass range, respectively, from 36 to 91% and 12 to 89% of that predicted in the absence of fishing (Table 1 and fig. S1). The variability in biomass over time and among stocks cannot be attributed entirely to fishing. Each stock has a unique recruitment history, some with periods of several years during which recruitment is more than 1 SD above or below the long-term average for the stock (fig. S2).

¹Joint Institute for Marine and Atmospheric Research, University of Hawaii, Honolulu, HI 96822, USA. ²Oceanic Fisheries Programme, Secretariat of the Pacific Community, BP D5, Noumea 98848, New Caledonia. ³Pacific Islands Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2570 Dole Street, Honolulu, HI 96822, USA. ⁴Inter-American Tropical Tuna Commission, 8604 La Jolla Shores Drive, La Jolla, CA 92037, USA.

*To whom correspondence should be addressed. E-mail: sibert@hawaii.edu

Longline fishing, which selectively removes the largest and oldest individuals, was the primary method of fishing for the first 25 years of the tuna fishery except in some coastal areas. The biomass of tunas larger than 175 cm fork length (measured from the tip of the snout to the center of the fork in the tail) decreased by 40% by the end of the 1970s (Fig. 3A). The current biomass of tunas larger than 175 cm is less than 17% of the biomass expected in the absence of fishing, but this segment of the population never

made up more than 5% of the total tuna biomass (fig. S5). The burgeoning purse-seine fishery in the 1980s began to remove smaller fish (~75 cm in length), spreading exploitation to earlier life-history stages (Fig. 3A).

An ontogenetic model relating size to trophic level was applied to the size structure of the catch, exploited population, and unexploited population to estimate trophic levels (2). The trophic level of the catch, aggregated across the eight stocks, was about 10% higher than the trophic

level in both the exploited and unexploited population in the 1950s and declined steadily to the same level as the exploited population (Fig. 3B). The trophic level of the catch dropped from 4.1 to 4.0 over the past 50 years because of the increased catch of smaller fish, but the trophic level in the exploited population has remained relatively constant at 3.9 (table S3).

Stock assessment is sometimes criticized (4) for dependence on unproven assumptions and for producing uncertain results. Integrated stock assessment models are, however, the tools of choice when attempting to accommodate disparate fisheries data as opposed to selecting a single data source [for example, (5)]. Our analysis depends mainly on ratios of estimated variables, which are less variable than point estimates (6). The analysis, which used all available fisheries data, shows that the biomass of top-level predators has not declined catastrophically in spite of the continuous increase in catches since the 1950s. The biomass of four stocks is greater than 74% of their unexploited potential, and the biomass of the remaining four stocks is 36 to 49% of the unexploited potential, as expected when the catch approaches MSY.

The biomass of several stocks of primary tuna and other top predators (7–10) has declined to MSY-associated levels. The ecosystem impact of such reductions is unknown, but model studies have shown that fishing all species in an ecosystem at mortality rates yielding single-species MSY may, under some conditions, lead to the erosion of trophic structure and depensa-





Fig. 1. Catches of major tuna species in the Pacific Ocean from 1950 through 2004 (1).

Fig. 2. Trends in total biomass for eight stocks of large predators in the Pacific Ocean, estimated from integrated stock assessment models. Blue lines indicate the biomass estimated from the observed fishing history (the exploited population), and red lines indicate the biomass estimated in the absence of all fishing (the unexploited population). The single black dash indicates the equilibrium biomass corresponding to MSY conditions, assuming current levels of recruitment and distribution of fishing mortality among fisheries. WCPO, western central Pacific Ocean; EPO, eastern Pacific Ocean.



tory effects on recruitment (11, 12). MSY is, nevertheless, enshrined by many national and international fishery management agencies as the target level of catch to which fishery management aspires. More conservative, but ad hoc, reference points, such as maintaining adult populations above 50% of the unexploited adult biomass, may be more appropriate for an ecosystem approach to fisheries management (13).

Attempting to establish an unvarying initial reference point or baseline against which to evaluate changes in biomass is misleading. For example, the biomass of yellowfin in the eastern Pacific is currently about 2 times greater than the biomass at the beginning of the data time series, suggesting that the stock is in good condition. The exploited biomass is, however, only 49% of the unexploited biomass, and the current catch is near MSY (14). Expressing the impact of fishing as the ratio of exploited to unexploited biomass is a more sensitive indicator than comparison of current biomass to the biomass at some arbitrary date in the past at which the stock is assumed to

be in a “pristine” or “virgin” state. The impact of the fishery is detectable in all stocks when the exploited-to-unexploited ratio is used.

The increase in the biomass of certain species, such as blue shark and skipjack, is a potentially important ecosystem response predicted by simple ecosystem models (15) and possibly attributable to a reduction in the biomass of other large predators. The possibility that the biomass of rapidly growing predators (such as *Coryphaena hippurus* and *Acanthocybium solandri*) may be also increasing should be examined. Unfortunately, complete fisheries statistics for these species are not available, and they rank low on the list of priority species for stock assessment in fishery management organizations.

Single-species assessment models do not explicitly include the effects of changes in the abundance of one species on the abundance of another. Until assessment models are able to simultaneously analyze multiple species, whether reduced abundance of larger fish has led to an increase in skipjack abundance will remain uncer-

tain. In spite of this limitation, single-species stock assessments have an important role in the ecosystem approach to fisheries. Single-species assessment methods applied to nontarget species, such as blue shark, reveal potentially important ecosystem trends.

The results presented here appear to differ sharply from widely accepted views of the status of large oceanic predatory fish stocks and the ecosystem effects of fishing (5, 16). Our results, however, should be interpreted as extensions of previous work, providing a more realistic (17–22) appraisal of the effects of fishing on the pelagic ecosystem. Our estimates of biomass trends interpret all available data from all major fishing fleets in the context of well-understood population dynamics processes. The analysis extends the concept of aggregating abundance across species (5) by summing biomass according to length across species, concluding that the biomass of fish smaller than 175 cm is near that predicted for stocks at full exploitation and that drastic declines in abundance are detectable only in fish larger than 175 cm. Furthermore, the analysis extends the concept of “shifting baseline” (23, 24) by defining an empirical, quantitative reference base that depends directly on the activity being analyzed, concluding that although some predator populations have declined severely in response to fishing, others have increased. Finally, the analysis extends the notion of examining the impact of fisheries on the trophic level of the catch (16) by also examining the trophic level of the population at large, concluding that there is no impact on the trophic level of the population and that the apparent reduction in the trophic level of the catch is caused by the development of purse-seine fisheries targeting smaller tunas; that is, by “fishing through the food web” (21).

Fishery scientists have been warning of impending stock conservation problems in Pacific yellowfin and bigeye tuna in the western and central Pacific Ocean since 2001 (25), and the relatively depressed condition of the spawning populations of these two species has stimulated fishery managers to attempt to impose constraints on the fishery. The Western and Central Pacific Fisheries Commission and the Inter-American Tropical Tuna Commission have the responsibility to implement conservation regulations in the western and eastern Pacific Ocean, respectively. Scientists in both commissions have recommended management options, including catch and effort limits, restrictions on the use of artificial floating objects by the purse-seine fishery, and time and area closures, to maintain sustainable fisheries for these species on both sides of the Pacific (26, 27). Unfortunately, these options have not been fully implemented, and decreases in fishing mortality have not yet been realized. Commissioners and national delegations to these commissions need to implement effective conservation measures before stocks reach a state where draconian measures, such as complete closures, are required to preserve the fishery and sustain the ecosystem.

Table 1. Comparison of estimated impacts of fishing computed as the ratio of estimated exploited biomass to the estimated unexploited biomass in the last year of the analysis (exploited/unexploited) and as the ratio of estimated exploited biomass in the last year to that in the first year (biomass ratio).

Stock	Exploited/unexploited		Biomass ratio	
	Total	Adult	Total	Adult
North Pacific albacore	0.38	0.12	1.34	0.94
Pacific bigeye	0.39	0.26	0.28	0.19
WCPO yellowfin	0.36	0.27	0.37	0.30
EPO yellowfin	0.49	0.41	2.12	2.27
North Pacific blue shark	0.91	0.89	1.20	1.16
WCPO skipjack	0.74	0.70	2.08	1.84
EPO skipjack	0.79	0.77	2.79	3.12
South Pacific albacore	0.89	0.83	0.89	0.91




Fig. 3. Two ecosystem indicators of the effects of fishing. (A) Impact of fishery on size. The ratio of exploited to unexploited biomass at increasing fork lengths by decade from 1950 through 2004 for all tuna stocks is shown. The dashed box indicates the size range of skipjack caught by the purse-seine fishery. The vertical dashed line indicates 175 cm fork length. (B) Impact of fishery on the trophic level of exploited tuna biomass and catch. The blue line is the ratio of the trophic level of exploited biomass to the trophic level of the unexploited biomass, and the green line is the ratio of the trophic level of the catch to the trophic level of the unexploited biomass. The dashed red line indicates 1.0, the relative trophic level in the unexploited biomass.

References and Notes

1. T. Lawson, Ed., *Western And Central Pacific Fisheries Commission Tuna Fishery Yearbook* (Western and Central Pacific Fisheries Commission, Pohnpei, Federated States of Micronesia, 2004), available at www.spc.int/oceanfish/Docs/Statistics/TYB.htm.
2. Background material and analytical methods are available as supporting material on *Science Online*.
3. The "exploited" biomass is defined here as the biomass of the population estimated by the assessment model. The "unexploited" biomass is defined as the biomass that might have existed in the absence of fishing, estimated by setting the fishing mortality parameters to zero in the model. The unexploited population trajectories include the estimated effects of temporal variability in recruitment on the dynamics of the populations, thus incorporating environmental constraints that occurred during the history of the fishery and the estimated effect of the additional recruitment resulting from a larger spawning biomass (2).
4. A. Longhurst, *Fish. Res.* **81**, 107 (2006).
5. R. Myers, B. Worm, *Nature* **423**, 280 (2003).
6. M. Labelle, *Fish. Res.* **71**, 311 (2005).
7. P. Kleiber, M. Hinton, Y. Uozumi, *Mar. Freshw. Res.* **54**, 349 (2003).
8. M. Labelle, NOAA Technical Memorandum *NOAA-TM-NMFS-SWFSC-341* (U.S. Department of Commerce, Washington, DC, 2002).
9. M. Hinton, W. Bayliff, *Inter-American Tropical Tuna Commission Stock Assessment Report* **3**, 328 (2002).
10. W. Bayliff, *Inter-American Tropical Tuna Commission Stock Assessment Report* **2**, 247 (2002).
11. C. Walters, V. Christensen, S. Martell, J. Kitchell, *ICES J. Mar. Sci.* **62**, 558 (2005).
12. S. Hall, J. Collie, D. Duplisea, M. Bravington, J. Link, *Can. J. Fish. Aquat. Sci.* **63**, 1344 (2006).
13. C. Walters, J. Kitchell, *Can. J. Fish. Aquat. Sci.* **58**, 39 (2001).
14. S. Hoyle, M. Maunder, *Inter-American Tropical Tuna Commission Stock Assessment Report* **6**, 3 (2005).
15. J. Hinke *et al.*, *Ecol. Soc.* **9**, 10 (2004).
16. D. Pauly, V. Christensen, J. Dalsgaard, R. Froese, F. Torres, *Science* **279**, 860 (1998).
17. J. Hampton, J. Sibert, P. Kleiber, M. Maunder, S. Harley, *Nature* **434**, E1 (2005).
18. T. Polacheck, *Mar. Policy* **30**, 470 (2006).
19. G. Burgess *et al.*, *Fisheries* **30**, 206 (2005).
20. C. Walters, *Can. J. Fish. Aquat. Sci.* **60**, 1433 (2003).
21. T. Essington, A. Beaudreau, J. Wiedenmann, *Proc. Natl. Acad. Sci. U.S.A.* **103**, 3171 (2006).
22. M. Maunder *et al.*, *ICES J. Mar. Sci.* **63**, 1373 (2006).
23. D. Pauly, *Trends Ecol. Evol.* **10**, 430 (1995).
24. J. Jackson *et al.*, *Science* **293**, 629 (2001).
25. *Report of the Fourteenth Meeting of the Standing Committee on Tuna and Billfish* (Secretariat of the Pacific Community, Noumea, New Caledonia, 2001), available at www.spc.int/oceanfish/Html/SCTB/SCTB14/sctb14rep.pdf.
26. J. Hampton *et al.*, *Estimates of Sustainable Catch and Effort Levels for Target Species and the Impacts on Stocks of Potential Management Measures* (WCPFC-SC1 Working Paper SA-WP-10, Western and Central Pacific Fisheries Commission, Pohnpei, Federated States of Micronesia, 2005), available at http://wcpfc.org/sc1/pdf/SC1_SA_WP_10.pdf.
27. M. Maunder, S. Harley, *Bull. Mar. Sci.* **78**, 593 (2006).
28. We thank our colleagues P. Crone, S. Hoyle, A. Langley, M. Ogura, Y. Takeuchi, M. Ichinokawa, and P. Williams, who assembled the data and conducted the stock assessments on which our synthesis is based; B. Bayliff, H. Dewar, G. Dinardo, K. Holland, K. Mason, C. Walters, G. Watters, and three anonymous reviewers for constructive comments on previous drafts of this paper; and D. Fournier for his work in creating the MULTIFAN-CL stock assessment method. This work was funded by Cooperative Agreement NA17R1230 between the Joint Institute for Marine and Atmospheric Research and the National Oceanic and Atmospheric Administration, by members of the Secretariat of the Pacific Community and its donor agencies, and by members of the Inter-American Tropical Tuna Commission. Views expressed in the paper do not necessarily represent the views of these agencies, their subagencies, or their member countries.

Supporting Online Material

www.sciencemag.org/cgi/content/full/314/5806/1773/DC1

Materials and Methods

SOM Text

Figs. S1 to S7

Tables S1 to S3

References and Notes

20 September 2006; accepted 8 November 2006

10.1126/science.1135347